

Quantification of large-scale and long-term groundwater resources in karst aquifers under Mediterranean climate: deterministic versus stochastic approaches

Flann Corcoran¹, **Simone Höglauer**¹, Lysander Bresinsky², Martin Sauter², Philippe Renard³, Edoardo Bucchignani⁴, Irina Engelhardt¹

¹Department of Hydrogeology, TU Berlin, Germany; ²University of Göttingen, Dept. of Applied Geology, Germany;

³University of Neuchâtel, Centre of Hydrogeology and Geothermics, Switzerland; ⁴Italian Aerospace Research Centre (CIRA), Capua, Italy.



Motivation

Water scarcity in many regions of the world will be exacerbated by **climate change**. **Carbonate aquifers** provide valuable water resource in the Mediterranean region, but are **vulnerable** to over-exploitation due to their low storage capacity.

Sustainable management of Mediterranean karst aquifers is a key issue at local & regional scale. However, the response of carbonate aquifers to **high-intensity precipitation events & droughts** is controlled by the distribution and type of karst features and the karst system.

Key Objective:

To **identify optimal modelling concepts** for highly dynamic & complex carbonate systems to improve management concepts for local water user management.

Study Area: Western Mountain Aquifer (WMA)

Located in Israel & Palestinian Territories (Figure 1).




Figure 1: WMA location (data from Abusaada & Sauter, 2012)

Cretaceous carbonate aquifer with developed karst system. Two limestone and dolomite sub-aquifers (both ~350 m thick) separated by argillaceous aquitard (~100 m thick).

Predicted Climate Change in the Recharge Area of the WMA (CIRA) (Figure 2).

- >2° increase in temperature.
- 20% reduction in precipitation.
- Reduction in frequency of very wet years.




Figure 2: Simulated Precipitation and Temperature in Recharge Areas of the WMA

Selection of a Suitable Numerical Modelling Approach

Two modelling approaches are used for comparison:

Deterministic multi-continuum approach

Model code: **HydroGeoSphere**.

- Considers surface routing, **unsaturated & saturated flow**.
- Recharge is simulated directly**.
- Parameterised with literature values
- Calibrated with piezometric pressure head & discharge time series.
- Enhanced simulation of response to extreme rainfall events.
- Subject to considerable **parameter uncertainty**.

Stochastic single-continuum approach

Model code: **MODFLOW & SKS**.

- Considers **saturated flow only**.
- Recharge from external calculation
- Parameterised with karstic networks from pseudo-genetic Stochastic Karst Simulator (SKS)
- Less reliant** on high-quality observation time-series.
- Stochastic parameterisation allows **estimation of uncertainty**.
- Fewer parameters, accounting for **data scarcity**.

Stochastic single-continuum workflow

Karstic networks will be generated using a Stochastic Karst Simulator (SKS) (Borghi et al.) for parameterisation of the single-continuum model (Figure 3).





Figure 3: Schematic flow diagram for stochastic single-continuum modelling approach

Deterministic multi-continuum model

Geological layers adapted from Abusaada & Sauter (2012) (Figures 4 and 5).

High- & low hydraulic conductivity domains used to represent conduits & fissured matrix, respectively.

First-order exchange between surface and sub-surface domains allows simulation of rapid and diffuse recharge.




Figure 4: E-W transect of multi-continuum model




Figure 5: Multi-continuum model discretisation and boundary conditions

First Results

Derived geological and climatic development (Figure 6)

Messinian Salinity Crisis (5.96 – 5.33 Ma) key period for karstification due to major sea-level decline & development of canyons at coast, allowing for development of karst to substantial depth.




Figure 6: Global and Mediterranean sea levels and climate conditions for Israel region. Data compiled from Miller et al. (2005), Vaks et al., (2013) and Frumkin et al. (2000).

Calibration Results for Multi-continuum Model

Calibrated to piezometric pressure head and discharge observations from Abusaada & Sauter (2012) (Figure 7).




Figure 7: a) Transient piezometric head at well Beteh Tiqva (coastal region); b) Simulated Vs observed heads for initial steady state run; c) Transient spring discharge at two spring locations

References

- Abusaada, M., Sauter, M., 2012. Studying the Flow Dynamics of a Karst Aquifer System with an Equivalent Porous Medium Model. Ground Water, 51(4), 641-50.
- Borghi, A., Renard, P. & Jenni, S., 2012. A pseudo-genetic stochastic model to generate karstic networks. Journal of Hydrology, 414-415, 516-529.
- Frumkin, A., Ford, D. C. & Schwarcz, H. P., 2000. Paleoclimate and vegetation of the last glacial cycles in Jerusalem from a speleothem record. Global Biogeochemical Cycles, 14.
- Laskow, M., Gendler, M., Goldberg, I., Gvirtzman, H. & Frumkin, A., 2011. 'Deep confined karst detection, analysis and paleo-hydrology reconstruction at a basin-wide scale using new geophysical interpretation of borehole logs', Journal of Hydrology 406(3), 158 – 169.
- Miller, K. G., Komiz, M. A., Browning, J. V., Wright, J. D., Mountain, G. S., Katz, M. E., Sugarman, P. J., Cramer, B. S., Christie-Blick, N. & Pekar, S. F., 2005. The Phanerozoic record of global sea-level change. Science, 310, 1293-1298
- Vaks, A., Woodhead, J., Bar-Matthews, M., Ayalon, A., Cliff, R. A., Zilberman, T., Matthews, A. & Frumkin, A., 2013. Pliocene–Pleistocene climate of the northern margin of Saharan–Arabian Desert recorded in speleothems from the Negev Desert, Israel. Earth and Planetary Science Letters, 368, 88-100.



Financing
This work is funded by the Federal Ministry of Education & Research (BMBF) of Germany

Contact
Technische Universität Berlin
Sekt. BH 3-2
Ernst-Reuter-Platz 1
10587 Berlin
simone.hoeglauer@tu-berlin.de
Tel: +49 30 314-72652